

ADULT AND PAEDIATRIC ENT HEARING AND BALANCE SPECIALIST

Central Sydney ENT
Suite 1 / 16 – 22 Burwood
Road, Burwood

Phone: (02) 9747-3199 Fax: (02) 9747-1499 Sydney ENT Hearing and Balance Centre

Suite 210, San Clinic Tulloch, 185 Fox Valley Road, Wahroonga

Phone: (02) 9989-8080 Fax: (02) 9475-0220

A. Prof Payal Mukherjee

MBBS, FRACS (ORLHNS), MS (USyd)

entcaresydney.com.au

POST OPERATIVE PROTOCOL OPERATION: NASAL SURGERY

HOSPITAL STAY: You may stay overnight or be discharged on the same day depending on the extent of surgery.

LEVEL OF ACTIVITY: A low level of activity is recommended. Avoid sport or strenuous activity for 2-3 weeks after the operation. Heavy exertion may cause post-operative bleeding.

- > It is normal to feel symptoms that mimic a flu.
- A poor level of energy is common.
- ➤ Nasal congestion and a significant amount of blood clot, bloody nasal discharge and post-nasal discharge is common. Your nose will feel blocked for the first 1-2 weeks due to inflammation, mucous and blood clots post surgery.
- This will decrease week 3 to week 4 after surgery. During this time breathe through your mouth and keep hydrated.
- > You will need to avoid blowing or rubbing your nose for the first 3-4 days, if you have to, blow your nose gently.
- > You will need to stay at home for 1 to 2 weeks

DIET: A normal diet is recommended. Avoid hot drinks and food for 3 days after the operation.

MEDICATIONS:

- You will be given post-operative antibiotics for 5 days to prevent an infection.
- You will be given some ointment to use in your nose if you have had a Septoplasty. You will need to apply this 3 times a day for 2 weeks to each side of your nose.

NASAL SPRAYS:

➤ Drixine or Oxymetazoline will be provided to be used predominantly to treat any mild to moderate postoperative bleeding. Should postoperative bleeding occur from 1 or both nostrils spray 3 to 5 sprays in each nostril to help control bleeding. Please call us if bleeding persists or is uncontrollable.

SALINE IRRIGATION:

- Should be used regularly during the entire recovery.
- The saltwater irrigation (FESS or FLO) is to be used liberally in both nostrils for 4-6 times a day in the first 2 weeks.
- > It may be cut down to 2 to 3 times a day thereafter depending on your recovery.



ADULT AND PAEDIATRIC ENT HEARING AND BALANCE SPECIALIST

Suite 1 / 16 – 22 Burwood Road, Burwood

Central Sydney ENT

Phone: (02) 9747-3199 Fax: (02) 9747-1499 Sydney ENT Hearing and Balance Centre

Suite 210, San Clinic Tulloch, 185 Fox Valley Road, Wahroonga

Phone: (02) 9989-8080 Fax: (02) 9475-0220

A. Prof Payal Mukherjee

MBBS, FRACS (ORLHNS), MS (USyd)

entcaresydney.com.au

NASAL PACKING:

- ➤ Generally there is dissolvable packing left inside your nose. This will not need to be removed but will dissolve or come out with the saline irrigation.
- Occasionally there may be more significant packing required and you will be notified should this be the case.

POST OP REVIEW: You will need to see your surgeon 3 weeks after surgery.

Please call the rooms if you have any questions after surgery.